


In the claims:

Cancel Claim 1 to 25.


Claim 26 (new) A method of treating Parkinson disease in warm-blooded animals comprising administering to warm-blooded animals in need thereof an amount of a compound of the formula



in racemic, enantiomeric form or any combination of these forms,

in which

A is



in which Q is selected from the group consisting of H, -OR²², -NR²³R²⁴, phenyl

optionally substituted by at least one substituent independently selected from the group

consisting of OH, cyano, nitro, alkyl, alkoxy or $\text{-NR}^{10}\text{R}^{11}$ and a group with two substituents representing together a methylenedioxy or ethylenedioxy, or Q is selected from the group consisting of -COPh , $\text{-SO}_2\text{Ph}$ or $\text{-CH}_2\text{Ph}$, said COPh , $\text{-SO}_2\text{Ph}$ or $\text{-CH}_2\text{Ph}$ optionally substituted on its aromatic parts by at least one independently alkyl or alkoxy or halogen,

R^{10} and R^{11} are independently selected from the group consisting of hydrogen, alkyl and -COR^{12} , or R^{10} and R^{11} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being independently selected from the group consisting of O, N and S,

R^{12} is selected from the group consisting of hydrogen, alkyl, alkoxy and $\text{NR}^{13}\text{R}^{14}$,

R^{13} and R^{14} are independently, hydrogen or alkyl, or R^{13} and R^{14} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being chosen independently from the group consisting of O, N and S,

R^{22} is selected from the group consisting of hydrogen, alkyl and aryl optionally substituted by at least one substituent selected from the group consisting of alkyl, OH, halogen, nitro and alkoxy,

R^{23} and R^{24} are independently selected from the group consisting of hydrogen, alkyl and -CO-R^{25} ,

R^{25} is alkyl,

and R^{19} , R^{20} and R^{21} are independently selected from the group consisting of hydrogen, halogen, -OH , -SR^{26} , alkyl, cycloalkyl, alkenyl, alkoxy, cyano, nitro, $\text{-SO}_2\text{NHR}^{49}$,

$-\text{CONHR}^{55}$, $-\text{S(O)}_q\text{R}^{56}$, $-\text{NH(CO)R}^{57}$, $-\text{CF}_3$, $-\text{OCF}_3$ and $\text{NR}^{27}\text{R}^{28}$,

R^{27} and R^{28} are independently selected from the group consisting of hydrogen, alkyl and $-\text{COR}^{29}$, or R^{27} and R^{28} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being independently selected from the group consisting of O, N and S,

R^{49} and R^{55} are independently each time that they occur, hydrogen or alkyl or alkylcarbonyl,

q is an integer from 0 to 2,

R^{56} and R^{57} are, each time that they occur, hydrogen or alkyl or alkoxy,

R^{29} is selected from the group consisting of hydrogen, alkyl, alkoxy and $-\text{NR}^{30}\text{R}^{31}$,

R^{30} and R^{31} are independently selected from the group consisting of hydrogen and alkyl, or R^{30} and R^{31} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being independently selected from the group consisting of O, N and S,

Y is O or S;

R^1 is hydrogen, alkyl, aminoalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, trifluoromethylalkyl, alkenyl, allenyl, allenylalkyl, alkynyl, cyanoalkyl, $-(\text{CH}_2)_g\text{Z}^1\text{R}^{39}$, $-(\text{CH}_2)_g\text{COR}^{40}$, $-(\text{CH}_2)_g\text{NHCOR}^{70}$, aryl, aralkyl, arylcarbonyl, heteroarylalkyl and aralkylcarbonyl, the aryl group of the aryl, aralkyl, arylcarbonyl,

heteroarylalkyl or aralkylcarbonyl itself being optionally substituted by at least one substituent selected from the group consisting of alkyl, halogen, alkoxy, nitro, cyano, cyanoalkyl, amino, alkylamino, dialkylamino, $-(CH_2)_k-Z^2R^{39}$ and $-(CH_2)_k-COR^{40}$, Z^1 and Z^2 is selected from the group consisting of a bond, -O-, $-NR^{41}$ - and -S-,

R^{39} and R^{41} are independently each time that they occur, selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl and cyanoalkyl,

R^{40} is, independently each time that it occurs, selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl, cyanoalkyl, alkoxy and $NR^{42}R^{43}$,

R^{42} and R^{43} are independently, each time that they occur, selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl and cyanoalkyl and R^2 is selected from the group consisting of hydrogen, alkyl, aminoalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, trifluoromethylalkyl,

$-(CH_2)_g-NHCOR^{71}$, aralkyl and heteroarylalkyl optionally substituted on the aryl or heteroaryl by at least one member selected from the group consisting of halogen, alkyl, alkoxy, hydroxyl, cyano, nitro, amino, alkylamino and dialkylamino,

R^{70} and R^{71} are independently alkyl or alkoxy;

or R^1 and R^2 , taken together with the carbon atom which carries them, form a carbocycle with 3 to 7 members;

B is selected from the group consisting of hydrogen, alkyl, $-(CH_2)_g-Z^3R^{44}$ and carbocyclic aryl optionally substituted 1 to 3 times by a member selected from the group

consisting of halogen, alkyl and alkoxy of 1 to 6 carbon atoms, hydroxy, cyano, nitro, amino, alkylamino, dialkylamino and a carbocyclic aryl,

Z^3 is selected from the group consisting of a bond, -O-, NR^{45} - and -S-,

R^{44} and R^{45} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, allenyl, allenylalkyl and cyanoalkyl;

Ω is OR^{48} ,

R^{50} and R^{52} are independently each time that they occur, selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, allenyl, allenylalkyl and cyanoalkyl, R^{51} is, independently each time that they occur, selected from the group consisting of hydrogen, cycloalkyl and cycloalkylalkyl in which the cycloalkyl has 3 to 7 carbon atoms, alkyl of 1 to 8 carbon atoms, alkenyl, alkynyl, allenyl, allenylalkyl, cyanoalkyl, alkoxyalkyl, $NR^{58}R^{59}$, aryl or aralkyl, said aryl or aralkyl optionally substituted by at least one member selected from the group consisting of halogen, alkyl and alkoxy,

R^{58} and R^{59} are independently a member selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, allenyl, allenylalkyl and cyanoalkyl,

R^{53} and R^{54} are independently a member selected from the group consisting of hydrogen or $-(CH_2)_k-Z^7R^{60}$ and $-(CH_2)_k-COR^{61}$,

Z^7 is a member selected from the group consisting of a bond, -O-, NR^{62} - and -S-,

R^{60} and R^{62} are independently a member selected from the group consisting of hydrogen, alkyl, alkenyl, allenyl, allenylalkyl, alkynyl, cyanoalkyl, aryl, aralkyl, arylcarbonyl,

aralkylcarbonyl, pyridinyl, pyridinylalkyl or pyridinylcarbonyl, the aryl or pyridinyl of the aryl, aralkyl, arylcarbonyl, aralkylcarbonyl, pyridinyl, pyridinylalkyl or pyridinylcarbonyl being optionally substituted by at least one substituent selected from the group consisting of alkyl, halogen, nitro, alkoxy, cyano, cyanoalkyl, $-(CH_2)_k-Z^8R^{63}$ and $-(CH_2)_k-COR^{64}$,

R^{61} is a member selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl, cyanoalkyl, alkoxy and $NR^{65}R^{66}$,

R^{65} and R^{66} are independently selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl and cyanoalkyl,

Z^8 is a member selected from the group consisting of a bond, $-O-$, $-NR^{67}-$ and $-S-$,

R^{63} and R^{67} are independently a member selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl and cyanoalkyl,

R^{64} is a member of the group consisting of hydrogen, alkyl, allenylalkyl, alkenyl, alkynyl, cyanoalkyl, alkoxy and $NR^{68}R^{69}$,

R^{68} and R^{69} are independently a member selected from the group consisting of hydrogen, alkyl, allenyl, allenylalkyl, alkenyl, alkynyl cyanoalkyl,


and R^{48} is a member selected from the group consisting of hydrogen, alkyl, alkynyl and cyanoalkyl;

g and p, each time that they occur, being independently integers from 1 to 6, and k and n, each time that they occur, being independently integers from 0 to 6;

or a salt thereof sufficient to treat Parkinson disease.

Claim 27 (new) The method of Claim 26 wherein

A is



in which Q is selected from the group consisting of H, -OR²², -SR²², -NR²³R²⁴, phenyl optionally substituted by at least one member selected from the group consisting of halogen, OH, cyano, nitro, alkyl, alkoxy and -NR¹⁰R¹¹ and a group of two substituents together being methylenedioxy or ethylenedioxy, or Q is selected from the group consisting of -COPh, -OPh, -SPh, -SO₂Ph or -CH₂Ph, said -COPh, -OPh, -SPh, -SO₂Ph or -CH₂Ph being optionally substituted on its aromatic part by at least one member selected from the group consisting of alkyl, alkoxy and halogen, R¹⁰ and R¹¹ are independently selected from the group consisting of hydrogen, alkyl, or R¹⁰ and R¹¹ form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 ring members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being independently selected from the group consisting of O, N and S, R²² is selected from the group consisting of hydrogen, alkyl and aryl optionally substituted by at least one member selected from the group consisting of alkyl, OH, halogen, nitro and alkoxy, R²³ and R²⁴ are, independently, hydrogen, or alkyl or -CO-R²⁵,

R^{25} is alkyl,

and R^{19} , R^{20} and R^{21} are independently selected from the group consisting of hydrogen, halogen, OH, SR^{26} , alkyl, cycloalkyl, alkenyl, alkoxy, cyano, nitro, $-SO_2NHR^{49}$, $-CONHR^{55}$, $-S(O)_qR^{56}$, $-NH(CO)R^{57}$, $-CF_3$, $-OCF_3$ and $NR^{27}R^{28}$,

R^{26} is hydrogen or alkyl,

R^{27} and R^{28} are independently selected from the group consisting of hydrogen, alkyl and $-COR^{29}$, or R^{27} and R^{28} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 ring members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being selected from the group consisting of O, N and S,

R^{49} and R^{55} are, independently each time that they occur, hydrogen or alkyl or alkylcarbonyl,

q is an integer from 0 to 2,

R^{56} and R^{57} are, independently each time that they occur, hydrogen or alkyl or alkoxy,

R^{29} is selected from the group consisting of hydrogen, alkyl, alkoxy and $-NR^{30}R^{31}$,

R^{30} and R^{31} are independently selected from the group consisting of hydrogen, alkyl, or

R^{30} and R^{31} form together with the nitrogen atom an optionally substituted heterocycle

containing 4 to 7 ring members and 1 to 3 heteroatoms including the nitrogen atom


already present, the additional heteroatoms being selected from the group consisting of O,

N and S.

Claim 28 (new)

The method of Claim 26 wherein

A is



in which Q is $-\text{OR}^{22}$, $-\text{SR}^{22}$, $-\text{NR}^{23}\text{R}^{24}$, phenyl optionally substituted by at least one member selected from the group consisting of halogen, OH, cyano, nitro, alkyl, alkoxy and $-\text{NR}^{10}\text{R}^{11}$,

R^{10} and R^{11} are independently selected from the group consisting of hydrogen or alkyl, or R^{10} and R^{11} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 ring members and 1 to 3 heteroatoms including the nitrogen atom already present, the additional heteroatoms being selected from the group consisting of O, N and S,

R^{22} is selected from the group consisting of hydrogen, alkyl and aryl optionally substituted by at least one member selected from the group consisting of alkyl, OH, halogen, nitro and alkoxy,

R^{23} and R^{24} are, independently, hydrogen or alkyl,


and R^{19} , R^{20} and R^{21} are independently selected from the group consisting of hydrogen, halogen, OH, $-\text{SR}^{26}$, alkyl, alkenyl, alkoxy and $\text{NR}^{27}\text{R}^{28}$,

R^{26} is hydrogen or alkyl,

R^{27} and R^{28} are independently selected from the group consisting of hydrogen, alkyl, or R^{27} and R^{28} form together with the nitrogen atom an optionally substituted heterocycle containing 4 to 7 members and 1 to 3 heteroatoms including the nitrogen atom already

present, the additional heteroatoms being selected from the group consisting of O, N and S.

Claim 29 (new) The method of Claim 28 wherein



in which Q is selected from the group consisting of $-OR^{22}$, $-SR^{22}$ and phenyl substituted by OH and optionally by at least one member selected from the group consisting of halogen, OH, alkyl and alkoxy,

R^{22} is hydrogen or alkyl,


and R^{19} , R^{20} and R^{21} are independently selected from the group consisting of hydrogen, halogen, OH, SR^{26} , alkyl and alkoxy,

R^{26} is hydrogen or alkyl.

Claim 30 (new) The method of Claim 29 wherein


the compound corresponds to formula (I)₃ in which Y is O

A is



in which Q is OH, two of the R^{19} , R^{20} and R^{21} are selected from the group consisting of alkyl, alkoxy, alkylthio, amino, alkylamino and dialkylamino and the third is selected from the group consisting of hydrogen, alkyl, alkoxy, alkylthio, amino, alkylamino and dialkylamino, or Q is phenyl substituted by OH and at least one member selected from the group consisting of halogen, OH, alkyl, alkoxy and $-NR^{10}R^{11}$, R^{10} and R^{11} are independently hydrogen or alkyl.


Claim 31 (new) The method of Claim 30 wherein A is



in which Q is OH, two of the R^{19} , R^{20} and R^{21} are alkyl and the third is H,
or Q is phenyl substituted by OH and at least one alkyl.

Claim 32 (new) The method of Claim 26 wherein Y is O;

A is



in which Q is OH, two of the R^{19} , R^{20} and R^{21} are alkyl and the third is hydrogen,


or in which Q is phenyl substituted by OH and at least one alkyl;

B is hydrogen;

n is 0 or 1;

R^1 and R^2 both are hydrogen;

Claim 33 (new) The method of Claim 32, wherein the compound is selected from the group consisting of



- 4-[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-N-methyl-2-thiazolemethanamine;
 - 2-(((4-[3,5-di(tert-butyl)-4-hydroxyphenyl]-1,3-thiazol-2-yl)methyl)(methyl)amino)-acetonitrile;
 - 2,6-di(tert-butyl)-4-(2-(((2-hydroxyethyl)(methyl)amino)methyl)-1,3-thiazol-4-yl)phenol;
 - 4-(((4-(3,5-di(tert-butyl)-4-hydroxyphenyl)-1,3-thiazol-2-yl)methyl)(methyl)amino)-butanenitrile;
 - 2,6-di(tert-butyl)-4-(4-(2-[methyl(2-propynyl)amino]ethyl)-1,3-oxazol-2-yl)phenol;
 - 3-(((2-[2-(3,5-di(tert-butyl)-4-hydroxyphenyl)-1,3-oxazol-4-yl]ethyl)(methyl)amino)-propanenitrile;
 - 2,6-di(tert-butyl)-4-(4-[2-(1-piperazinyl)ethyl]-1,3-oxazol-2-yl)phenol;
- and the pharmaceutically acceptable salts of the latter.

Claim 34 (new) The method of Claim 26 wherein the compound is selected from the group consisting of 2-[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-4-oxazoleethanol, 2,6-ditert-butyl-4-(4-{2-[methyl(2-propynyl)amino]ethyl}-1,3-oxazol-2-yl)phenol, [{2-[2-(3,5-ditert-butyl-4-hydroxyphenyl)-1,3-oxazol-4-yl]ethyl}(methyl)amino]acetonitrile, 3-[{2-[2-(3,5-ditert-butyl-4-hydroxyphenyl)-1,3-oxazol-4-yl]ethyl}(methyl)amino]-propanenitrile, 2,6-ditert-butyl-4-{4-[2-(1-piperazinyl)ethyl]-1,3-oxazol-2-yl}phenol hydrochloride, 2,6-ditert-butyl-4-{4-(hydroxymethyl)-1,3-thiazol-2-yl}phenol and 2,6-ditert-butyl-4-{4-[(methylamino)methyl]-1,3-thiazol-2-yl}phenol hydrochloride.

Claim 35 (new) The method of Claim 34 wherein the disorder is selected from the group consisting of Parkinson's disease, Alzheimer's disease, Huntington's chorea, amyotrophic lateral sclerosis and peripheral neuropathies.